

เอกสารอ้างอิง

- โอภา วัชรคุปต์ และคณะ. **สารต้านอนุมูลอิสระ**. พิมพ์ครั้งที่ 1. กรุงเทพฯ : พี. เอส. พรินท์, 2006.
- กรมวิทยาศาสตร์บริการ กระทรวงวิทยาศาสตร์และเทคโนโลยี. แอนโทไซยานิน. 2010.
- Senevirathne M, Kim SH, Siriwardhana N, Ha JH, Lee KW, Jeon YJ. (2006). *Antioxidant potential of ecklonia cava on reactive oxygen species scavenging, metal chelating, reducing power and lipid peroxidation inhibition*. Food Science and Technology International. 12 (1): 27-38.
- Silva BM, Andrade PB, Valent P, Ferreres F, Seabra R M, Ferreira MA. (2004). *Quince (Cydonia oblonga Miller) fruit (pulp, peel, and seed) and jam: Antioxidant activity*. Journal of Agricultural and Food Chemistry. 52 (15): 4705-4712.
- Giasson BI, Ischiropoulos H, Lee VMY, Trojanowski JQ. (2002). *The relationship between oxidative/nitrative stress and pathological inclusions in Alzheimer's and Parkinson's diseases*. Free Radical Biology and Medicine. 32 (12), 1264-1275.
- Cheok CY, Chin NL, Yusof YA, Talib RA, Law CL. (2013). *Optimization of total monomeric anthocyanin (TMA) and total phenolic content (TPC) extractions from mangosteen (Garcinia mangostana Linn.) hull using ultrasonic treatments*. Industrial Crops and Products. 50, 1-7.
- Abdel-Hameed ESS. (2009). *Total phenolic contents and free radical scavenging activity of certain Egyptian Ficus species leaf samples*. Food Chemistry. 114 (4): 1271-1277.
- Chen LG, Yang LL, Wang CC. (2008). *Anti-inflammatory activity of mangostins from Garcinia mangostana*. Food Chem Toxicol. 46 (2): 688-693.
- Bouayed J, Bohn T. *Exogenous antioxidants - Double-edged swords in cellular redox state: Health beneficial effects at physiologic doses versus deleterious*

- effects at high doses.* Oxidative Medicine and Cellular Longevity. 3 (4): 228-237.
- Gan CY, Latiff AA. (2011). *Extraction of antioxidant pectic-polysaccharide from mangosteen (Garcinia mangostana) rind: Optimization using response surface methodology.* Carbohydrate Polymers. 83 (2): 600-607.
- Wang JJ, Shi QH, Zhang W, Sanderson BJ. (2012). *Anti-skin cancer properties of phenolic-rich extract from the pericarp of mangosteen (Garcinia mangostana Linn.).* Food Chem Toxicol. 50 (9): 3004-3013.
- Yu L, Zhao M, Yang B, Zhao Q, Jiang Y. (2007). *Phenolics from hull of Garcinia mangostana fruit and their antioxidant activities.* Food Chemistry. 104 (1): 176-181.
- A.O.A.C. (2000). *Official Methods of Analysis of AOAC International.* Gaithersburg, Md.: Association Official Analytical Chemists.
- Waterhouse A, Wiseman S, Korver O. (2001). *The health effects of tea and tea components: Opportunities for standardizing research methods.* Crit. Rev. Food. Sci. Nutr. 41: 387-412.
- Bushra S, Farooq A, Muhammad A. (2009). *Effect of Extraction Solvent/Technique on the Antioxidant Activity of Selected Medicinal Plant Extracts.* Molecules. 14: 2167-2180.
- Brand-Williams W, Cuvelier ME, Berset C. (1995). *Use of a free radical method to evaluate antioxidant activity.* Lebensmittel-Wissenschaft und Technologie. 28: 25-30.
- Rice-Evans CA, Miller NJ. (1996). *Antioxidant activities of flavonoids as bioactive components of food.* Biochem. Soc. Trans. 24: 790-795.