

## บรรณานุกรม

### บรรณานุกรมภาษาไทย

- ประสาท บริสุทธิ์เพ็ชร และคณะ. (2551). การทดสอบฤทธิ์ต้านเชื้อของสมุนไพรรักษาในห้วงปฏิบัติการ. *เอกสารประชุมวิชาการสัตวแพทย์ศาสตร์ มหาวิทยาลัย ขอนแก่น ครั้งที่ 9 “สัตวแพทย์ ทางเลือกวันนี้”* 11 - 12 มิถุนายน 2551 ณ ห้องประชุมใหญ่ คณะสัตวแพทย์ มหาวิทยาลัยขอนแก่น.
- พันธิวา กระจ่าง, ประพนอม จันทโรนัย และ พิมพวีดี พรพงศ์ รุ่งเรือง. (2013). การจำแนกพันธุ์ สมอ (*Terminalia chebula* Retz.) วงศ์ หูกวาง (Combretaceae). *KKU Research Journal*. 18 (6), 937-948.
- มหาวิทยาลัยอุบลราชธานี. คณะเภสัชศาสตร์, (มปป). “ฐานข้อมูลสมุนไพรไทย” สืบค้นเมื่อ 2558, มิถุนายน 15, เข้าถึงได้จาก <http://www.phargarden.com/main.php>
- อรอนงค์ พริ้งศุลกะ. (2555). *จุลชีววิทยาทางการแพทย์: แบคทีเรียก่อโรค*. กรุงเทพมหานคร: จรัส สนิทวงศ์การพิมพ์.
- อรัญญา มโนสร้อย และ จีระเดช มโนสร้อย. (2550). *ไลโปโซมสำหรับยาผ่านทางผิวหนังและ เครื่องสำอาง*. พิมพ์ครั้งที่ 1. กรุงเทพมหานคร: โอ.เอส. พรินติ้งเฮาส์.

### บรรณานุกรมภาษาต่างประเทศ

- Ahmed, I., & Beg, AZ. (2001). Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *Journal of Ethnopharmacology*, 74, 113-123.
- Ahmad I, Mehmood Z ,& Mohammad F (1998). Screening of some Indian medicinal plants for their antimicrobial properties. *Journal of Ethnopharmacology*. 62(2), 183-193
- Aneja, K.R., & Joshi, R. (2009). Evaluation of antimicrobial properties of fruit extracts of *Terminalia chebula* against dental caries pathogens. *Jundishapur Journal of Microbiology*, 2, (3) 105-111.
- Azimi,H., Fallah-Tafti, M., Khakshur, A.A., & Abdollahi, M. (2012). A review of phytotherapy of acne vulgaris : Perspective of new pharmacological treatments. *Fitoterapia*, 83,1306-1317.
- Baumann, L., & Keri, J. Acne (Type 1 Sensitive Skin) In: Baumann, L., Saghari, S., & Weisberg, s. (eds). *Cosmetic Dermatology Principles and Practice*. 2<sup>nd</sup> ed. New York: McGraw-Hill Companies, 2009, 121-127.

- Bag, A., Bhattacharyya, S.K., Bharati, P., Pal, N.K., & Chattopadhyay R.R. (2009). Evaluation of antibacterial properties of Chebulic myrobahan (fruit of *Terminalia chebula* Retz.) extract against methicillin resistant *Staphylococcus aureus* and trimethoprim-sulphamethoxazole resistant uropathogenic *Escherichia coli*. *African Journal of Plant Science*, 3, 025-029
- Bag, A., Bhattacharyya, S.K., & Chattopadhyay, R.R. (2013). The development of *Terminalia chebula* Retz. (Combretaceae) in clinical research. *Asian Pacific Journal Tropical Biomedicine*, 3 (3), 244-252
- Bag, A., Bhattacharyya, S.K., Pal, N.K., & Chattopadhyay, R.R. (2012). In vitro antimicrobial potential of *Terminalia chebula* fruit extracts against multidrug-resistant uropathogens. *Asian Pacific Journal of Tropical Biomedicine*, S1883-S1887.
- Barazani, V.O., Sathiyamoorthy, P., Shalev, R., Vardy, D., & Golan, G.A. (2003). Screening of South-Indian medicinal plants for anti-fungal activity. *Phytotherapy Research*, 17, (9) 1123-1125.
- Bonjar, G.H. (2004). Inhibition of Clotrimazole-resistant *Candida albicans* by plants used in Iranian folkloric medicine. *Fitoterapia*, 75, 74-76.
- Chattopadhyay, R.R., & Bhattacharyya, S.K. (2007). Plant review *Terminalia chebula*: An update. *Pharmacognosy Reviews*, 1, 151-156.
- Chiller, K., Selkin, B.A., & Murakawa, G.J. (2001). Skinmicroflora and bacterial infections of the skin. *Journal of Investigative Dermatology Symposium Proceedings*, 6, 170-174.
- Chomnawang, M.T., Surassmo, S., Nukoolkarn, V.S., & Gritsanapan, W. (2005). Antimicrobial effects of Thai medicinal plants against acne-inducing bacteria. *Journal of Ethnopharmacology*, 101, 330-333.
- Chrubasik, S., Pittler, H. M. & Roufogalis, B. D. (2005). Zingiberis rhizome : A comprehensive review on the ginger effect and efficacy profiles. *Phytomedicine*, 12, 684-701.
- Cogen, A.L., Nizet, V., & Gallo, R.L. (2008). Skinmicrobiota: a source of disease of defence. *British Journal of Dermatology* , 158, 442-455.
- Dash, B. (1991). *MateriaMedica of Ayurveda*. New Delhi: B. Jain Publishers, New Delhi, 170-174.
- Dixit, D., Dixit, A.K., Lad, H., Gupta, D., & Bhatnagar, D. (2013) Radioprotective effect of *Terminalia chebula* Retzius extract against g-irradiation-induced oxidative stress. *Biomedicine & Aging Pathology*, 3, 83-88.

- Dutta, B.K., Rahman, I., & Das, T.K. (1998). Antifungal activity of Indian plant extracts. *Mycoses*, 41, (11-12) 535-536.
- Eady, E.A., & Ingham, E. (1994). *Propionibacterium acnes*- friend or foe?. *Reviews in Medical Microbiology*, 5, 163-173
- Fierer, N., Lauber, C.L., Zhou, N., McDonald, D., Costello, E.K., & Knight, R. (2010). Forensic identification using skin bacterial communities. *Proceedings of the National Academy of Sciences USA*, 107, 6477-6481.
- Ghosh A., Das, B.K., Roy, A., Mandal, B., & Chanda, G. (2008). Antibacterial activity of some medicinal plant extracts. *Journal of Natural Medicines*, 62, 259-262.
- Grice, E.A., & Segre, J.A. (2011). The skin microbiome. *Nature Reviews Microbiology*, 9, 244-253.
- Jappe, U., Ingham, E., Henwood, J., & Holland, K.T. (2002). *Propionibacterium acnes* and inflammation in acne; *P. acnes* has T-cell mitogenic activity. *British Journal of Dermatology*, 146, 202-209.
- Kannan, P., Ramadevi, S.R., & Waheeta, H. (2009). Antibacterial activity of *Terminalia chebula* fruit extract. *African Journal of Microbiology Research*, 3 (4), 180-184
- Kathrivel, A., & Sujatha, V. (2012). *In vitro* assessment of antioxidant and antibacterial properties of *Terminalia chebula* Retz. *Leaves Asian Pacific Journal of Tropical Biomedicine*, S788-S795.
- Khan, K.H. (2009). The effect of regular intake of *Terminalia chebula* on oxidative stress in mice originated from *Salmonella typhimurium*. *EurAsian Journal of BioSciences*, 3, 113-121.
- Khan, K.H., & Jain, S.K. (2009). Regular intake of *Terminalia chebula* can reduce the risk of getting typhoid fever. *Advanced Biotech*, 8, 10-15.
- Kong, H.H. (2011). Skinmicrobiome: genomics-based insights into the diversity and role of skin microbes. *Trends in Molecular Medicine*, 17, 320-328.
- Leyden, J.J., McGinley, K.J., & Vowels, B. (1998). *Propionibacterium acnes* colonization in acne and non acne. *Dermatology*, 196, 55-58.
- Lodes, M.J., Secrist, H., Benson, D.R., Jen, S., Shanebeck, K.D., Guderian, J., Maisonneuve, J.F., Bhatia, A., Persing, D., Patrick, S., & Skeiky, Y.A.W. (2006). Variable expression of immunoreactive surface proteins of *Propionibacterium acnes*. *Microbiology*, 152, 3667-3681.

- Malckzadeh, F., Ehsanifar, H., Shahamat, N., Levin, M., & Colwell R.R. (2001). Antibacterial activity of black myrobalan (*Terminalia chebula* Retz.) against *Helicobacter pylori*. *International Journal of Antimicrobial Agents*, 18, 85-88.
- Marples, M. (1965). *The Ecology of the Human Skin*. Charles C Thomas, Bannerstone House.
- Ramgopal Mopuri, R., & Meriga B. (2014). In vitro anti oxidant activity and acute oral toxicity of *Terminalia paniculata* bark ethanolic extract on Sprague Dawley rats. *Asian Pacific journal of tropical biomedicine*, 4(4), 294-298.
- Naik, G.H., Priyadarsini, K.I., Naik, D.B., Gangabhairathi, R., & Mohan, H. (2004). Studies on the aqueous extract of *Terminalia chebula* as a potent antioxidant and a probable radioprotector. *Phytomedicine*, 11, 530-538.
- Nishijima, S., Kurokawa, I., Katoh, N., & Watanabe, K. (2000). The bacteriology of acne vulgaris and antimicrobial susceptibility of *Propionibacterium acnes* and *Staphylococcus epidermidis* isolated from acne lesions. *The Journal of Dermatology*, 27, 318-323.
- Noble, W.C. (2004). *The skin microflora and microbial diseases*. Philadelphia PA, Cambridge University Press.
- Noble, W.C., & Somerville, D.A. eds. (1974). *Microbiology of Human Skin*. W.B. Saunders Company Ltd.
- Pfundstein, B., Desouky, S. El K., Hull, W.E., Haubner, R., Erben, G., & Owen, R.W. (2010). Polyphenolic compounds in the fruits of Egyptian medicinal plants (*Terminalia bellerica*, *Terminalia chebula* and *Terminalia horrida*): Characterization, quantitation and determination of antioxidant capacities. *Phytochemistry*, 71, 1132-1148.
- Rangsrivong, P., Rangkadilok, N., Satayavivad, J., Goto, M., & Shotipruk, A. (2009). Subcritical water extraction of polyphenolic compounds from *Terminalia chebula* Retz. Fruits. *Separation and Purification Technology*, 66, 51-56.
- Sato, Y., Oketani, H., Singyouchi, K., Ohtsubo, T., Kihara, M., & Shibata, H. (1997). Extraction and purification of effective antimicrobial constituents of *Terminalia chebula* Retz. Against methicillin resistant *Staphylococcus aureus*. *Biological and Pharmaceutical Bulletin*, 20, 401-404
- Srivastava, P., Raut, H.N., Wagh, R.S., Puntambekar, H.M., & Kulkarni, M.J. (2012). Purification and characterization of an antioxidant protein (~16 kDa) from *Terminalia chebula* fruit. *Food Chemistry*, 131, 141-148.

- Surveswaran, S., Cai, Y.Z., Corke, H., & Sun, M. (2007). Systematic evaluation of natural phenolic antioxidants from 133 Indian medicinal plants. *Food Chemistry*, *102*(3), 938-953.
- Webster, G.F., Leyden, J.J., Norman, M.E., & Nilsson, U.R. (1978). Complement activation in acne vulgaris: in vitro studies with *Propionibacterium acnes* and *Propionibacterium granulosum*. *Infection and Immunity*, *22*, 523-529.
- White, G.M. (1991). Recent findings in the epidemiology, classification and subtypes of acne vulgaris. *Journal of the American Academy of Dermatology*, *39*, 534-537.
- Zvulunov, A. Acne In: Shai, A., Maibach, H.I., & Baran, R. (eds). *Handbook of Cosmetic Skin Care*. 2<sup>nd</sup> ed. London: Informa UK Ltd, 2009, 58-76.