



Assessing Non-Linearity and Stationarity in the Time Series of Albania's Annual Emissions of CO₂ from Land-Use Change

Orgeta Gjermëni*

*Department of Mathematics and Physics, Faculty of Technical and Natural Sciences,
University of Vlora "Ismail Qemali", Vlora 9400, Albania*

Received 28 July 2024; Received in revised form 23 November 2024

Accepted 3 December 2024; Available online 27 December 2024

ABSTRACT

The annual emissions of CO₂ from land-use change in Albania are the main focus of this research. The aim is to analyze the presence of non-linearity and stationarity. A mixed-methods strategy is used, which combines descriptive, inferential, and exploratory data analysis in time series data. A data sample was obtained from the Our World in Data website, spanning from 1850 through 2022. After the Isolation Forest technique was employed to identify outliers in the time series, the Long-Short-Term Memory model was used to impute them. Exploratory data analysis was applied to the original and imputed time series to ensure that the basic characteristics of the initial data distribution were preserved. Non-linearity and stationarity were checked in the imputed time series before and after applying the first differences. Non-linearity was assessed using the BDS test and the Teräsvirta Neural Network test. In the presence of non-linearity, stationarity was analyzed using the KPSS test, the Zivot-Andrews Unit Root test, and the Breitung test. The first differencing application transformed the non-stationary series into a stationary one, but it was insufficient to eliminate non-linearity. This highlights the complex nature of CO₂ emissions data and the need for sophisticated modeling techniques.

Keywords: Carbon emissions; Land-use impacts; LSTM imputation; Non-linearity; Stationarity tests

1. Introduction

Agriculture, forestry, and other land-use sectors are some of the contributors to the total greenhouse gas (GHG) emissions in Albania. The main emitters are the 'livestock' with 41%

of the total and the 'land' with 38% of the total GHG emissions. Although forests are believed to be a sink of GHG emissions, under the category 'land', they represent one of the key sources of emissions, mostly due to their

neglected management in recent years. Furthermore, negative impacts in this direction have arisen from uncontrolled deforestation, massive forest fires, a lack of effective investment in forest improvement and afforestation, informality, and an absence of development reforms [1].

In an empirical investigation [2] with data from Malaysia (1990–2019), it was shown that agricultural land expansion by 1% is associated with an increase of 0.84% in carbon dioxide (CO₂) emissions in the long run. Meanwhile, a 1% reduction in the wooded area, has resulted in a 5.41% long-term impact on higher CO₂ emissions.

Carbon dioxide emissions from land-use change (CO₂-E-LUC) are considered a summation of various activities that emit carbon stored in vegetation or soil [3, 4]. This includes emissions from: deforestation (CO₂-E_{def.}), forest degradation (CO₂-E_{deg.}), conversion of forest land to agriculture (CO₂-E_{conv.agric.}), conversion of natural land to urban areas (CO₂-E_{urb.conv.}), soil degradation, and wetland drainage (CO₂-E_{soil deg. and wetland drain.}). As expressed in a formula we have,

$$\begin{aligned} \text{CO}_2\text{-E-LUC} = & \text{CO}_2\text{-E}_{\text{def.}} + \text{CO}_2\text{-E}_{\text{deg.}} \\ & + \text{CO}_2\text{-E}_{\text{conv.agric.}} + \text{CO}_2\text{-E}_{\text{urb.conv.}} \\ & + \text{CO}_2\text{-E}_{\text{soil deg. and wetland drain.}} \end{aligned} \quad (1.1)$$

Trajectories of land-use change indicate positive and negative relationships between man and the environment. Land-use change analyses are necessary to assist the government in appropriate zoning to minimize or eliminate negative environmental impacts. With the beginning of land trading in the absence of regulatory legislation from 1996 to 2003 in Albania, land-use changes were more dynamic [5]. According to [6], land-use change caused 215 and 142 Pg C of global emissions and removals, respectively, between 1961 and 2020, resulting in an average net emissions of 1.21 Pg C per year.

The state of climate change in six Western Balkan countries (Albania included) was studied in [7]. The development of

sustainable forestry, the improvement of forest management practices, and the rehabilitation of degraded forest land are integral parts of the strategies that have to do with land-use change and forestry.

The annual emissions of CO₂ from land-use change in Albania, as measured in tonnes per person (CO₂-E-LUC per capita), are the primary focus of analysis in this research, seen from a statistical perspective. CO₂-E-LUC per capita is expressed by,

$$\text{CO}_2\text{-E-LUC per capita} = \frac{\text{CO}_2\text{-E-LUC}}{\text{population}} \quad (1.2)$$

There are several reasons, such as climate change [8, 9], health [10], economic development [11, 12], carbon budgets [13], etc., why it is important to conduct research on the CO₂-E-LUC per capita time series data in Albania as well as abroad. Further, to our knowledge, in Albania, there is a lack of studies in the area of CO₂ emissions data, divided by their categories.

Albania is committed to implementing policies to lower GHG emissions from various economic sectors. By December 2022, the country had partially aligned with the Regulation on the Governance of the Energy Union and Climate Action. The level of emission reductions planned for 2021–2030 in the National Energy and Climate Plan adopted in 2021 is 18.7%. The plan relies significantly on the reduction of CO₂ by forests, but in contrast with this, there are very limited financial means and capacity allocated to protecting and managing forests, including measures to promote reforestation and manage forest fire risks. In February 2023, a ministerial decision approved the National Strategy for Development and European Integration 2022–2030, which sets out a series of priority measures like adopting climate secondary legislation, climate budgeting, and nature-based solutions. Furthermore, Albania needs to address strategic investment planning, and the implementation and monitoring capacity of infrastructure projects [14].

The main objective of this research is to analyze the presence of non-linearity and stationarity in the time series data of annual CO₂-E-LUC per capita. To further study this objective, we raise the following research questions:

- RQ1: Is there non-linearity in the time series data?
- RQ2: Is the time series data stationary?

Non-linearity and non-stationarity are processes that are often encountered in time series, and they can be the result of complex underlying dynamics. If these issues are not addressed before building specific predictive models, it would compromise the reliability of the results. In the context of CO₂ emissions, since many decision-making policies are based on predictive models of time series of

emissions, they are directly influenced by the accuracy of these models.

2. Materials and Methods

The present research implemented a mixed-methods approach to investigate the presence of non-linearity and stationarity in the annual CO₂-E-LUC per capita series data. It is a combination of inferential, descriptive, and exploratory methods. Data sampling related to Albania’s annual CO₂-E-LUC per capita was gathered from Our World in Data, available for download at [15]. Ritchie in [16] provides sources and methods used to produce CO₂ emissions dataset. The data timeframe that is analyzed herein spans from 1850 to 2022. The variables under study are “annual CO₂-E-LUC per capita” and “year”, analyzed as a time series.

Table 1. The non-linearity and stationarity tests applied.

Purpose	Test	Function utilized in R for the test implementation	Null hypothesis	Alternative hypothesis	Rejection of Null hypothesis
Non-linearity testing RQ1	The Brock-Dechert-Scheinkman (BDS) Test [17, 18]	bds.tst() [19] with embedding dimensions values (2, 3) and four epsilon values (0.5SD, 1SD, 1.5SD, 2SD).	The time series is independently and identically distributed (i.i.d.).	The time series is not i.i.d. (exhibits nonlinearity or chaotic behavior).	p<0.05
	The Teräsvirta Neural Network Test [20, 21]	terasvirta.test() [22]	The time series follows a linear model.	The time series follows a nonlinear model.	p<0.05
Stationarity testing RQ2	The Kwiatkowski-Phillips-Schmidt-Shin (KPSS) Test [23]	ur.kpss() [24] type= c(“mu”, “tau”) lags=c(“short”, “long”) ¹	The time series has stationarity around a mean / deterministic trend.	The time series is not level / trend-stationary.	The test statistic exceeds the critical values (1%, 2.5%, 5%, 10%).
	Zivot-Andrews Unit Root Test [25]	ur.za() [24] model =c(“intercept”, “trend”, “both”) Optimal lag value ²	The time series has a unit root and is non-stationary, with no structural break.	The time series is stationary with a structural break at some unknown point in the time series.	The test statistic is less than the critical value values (1%, 5%, 10%)
	Breitung Test (Elliot, Rothenberg, and Stock Unit Root Test) [26, 27]	ur.ers() [24] type= “P-test” model= c(“constant”, “trend”)	The time series has a unit root, it is non-stationary around a constant mean/deterministic trend.	The time series is stationary around a constant mean/deterministic trend.	The test statistic is less than the critical values (1%, 5%, 10%)

Note 1: lags= “short” sets the number of lags to $4 \left(\frac{n}{100}\right)$, whereas lags= “long” sets the number of lags to $12 \left(\frac{n}{100}\right)$.

Note 2: The Akaike Information Criterion (AIC) [28] was used to select the optimal lag length, rotating from 1 to 40, minimizing the AIC value. This ensures that the model includes enough lags to capture autocorrelation without overfitting.

The statistical procedure started with the identification of time series outliers. Their presence and mishandling can affect

the accuracy of predictive models. The Isolation Forest technique [29, 30] was used to detect them. This technique effectively

identifies anomalies without assuming distributional properties, making it well-suited for non-linear, non-stationary time series data. Random binary trees from subsets of the data are constructed to isolate individual points based on their anomaly

scores. A range of trees (50, 100, 150, and 200) were evaluated, and the optimal number was selected using cross-validation. Outliers were identified as points that exceeded the 95th percentile threshold of the anomaly scores.




Fig. 1. Flowchart of statistical procedure steps.

After that, the outliers were replaced with NA values. The Long Short-Term Memory (LSTM) model [31-34] was chosen for the imputation, due to its ability to capture

complex temporal dependencies in non-linear and non-stationary data. The model architecture details were as follows: two layers, each containing 30 units; a specified

timestep of 5; 100 epochs; and a batch size 16. Model performance [35, 36] was evaluated using metrics: Root Mean Squared Error (RMSE), Mean Absolute Percentage Error (MAPE), Symmetric Mean Absolute Percentage Error (sMAPE), Mean Absolute Scaled Error (MASE), and R-squared (R^2).

Furthermore, it was continued with an exploratory data analysis of both original and imputed time series data to verify if the basic characteristics of the initial data distribution were preserved. The summary statistics used were: minimum (Min), first quartile (Q1), median (Q2), mean, third quartile (Q3), maximum (Max), standard deviation (SD), skewness, and kurtosis. The data was tested for normality using the Anderson-Darling normality test [37]. Locally estimated scatterplot smoothing (LOESS) was obtained through the function `loess()` with `span=0.3` [38] [39].

Graphical representations such as boxplots, line plots, density plots, and LOESS trend plots, were used to visually see the approximation made by the imputation method. Before checking for the stationarity of a series, it is necessary to analyze its linearity, because some tests that assess stationarity assume that the series is linear. Conversely, if the non-linearity of the series is confirmed, then the tests for stationarity should be chosen under this circumstance.

The applied tests to check on the two research questions are described with implementation details in Table 1. All the procedure steps in analyzing the time series data are outlined in the flowchart in Fig. 1. The statistical computations are gathered using R software [40].

3. Results and Discussion

3.1 Exploratory Data Analysis

The optimal number of trees chosen by cross-validation was 50, while applying the Isolation Forest technique. Nine outliers'

values were identified in the time series data of annual CO2-E-LUC per capita, using the threshold 0.6237087. These values represent only 5% of the sample. Together with the corresponding years and the imputed values, they are presented in Table 2.

Table 2. Outliers, corresponding imputed values, and the respective years.

Year	Outlier	Imputed
1951	3.536	3.171
1954	4.071	3.650
1955	4.163	3.569
1956	4.238	3.446
1957	4.348	3.302
1959	4.401	3.671
1960	3.755	3.632
2002	-0.113	0.048
2004	-0.312	0.048

Table 3. Summary statistics of time series data before and after imputation.

Statistic	Before	After
Min	-0.312	-0.071
Q1	1.060	1.060
Q2	2.56	2.56
Mean	2.08	2.06
Q3	2.891	2.891
Max	4.40	4.37
SD	1.17	1.12
Skewness	-0.499	-0.628
Kurtosis	-0.811	-0.93
Anderson-Darling Test	Test Statistic (A):	Test Statistic (A):
	8.3608	9.7374
	p-value:	p-value:
	< 2.2e-16	< 2.2e-16

The LSTM model imputation achieved an RMSE of 0.1336, indicating a minimal average error, and MAPE of 2.06%, emphasizing a low percentage of error. The sMAPE of 2.93%, further supported the consistency of the imputation. Additionally, the MASE of 0.279 showed scaled accuracy, while the R^2 of 0.988 confirmed a near-perfect fit between imputed and outliers' values.

Future analysis could be conducted to investigate what might have occurred in these years to result in these extreme values for annual CO2-E-LUC per capita, but this is not the focus of the current study.




Fig. 2. Box plots of annual CO2-E-LUC-per capita before and after imputation.




Fig. 3. Density plots of annual CO2-E-LUC-per capita before and after imputation.

In the following, the imputed time series is obtained from the original time series, where outlier values have been replaced by the imputed values. The summary statistics are presented in Table 3. The visual representations are given by box plots in Fig. 2 and density plots in Fig. 3. When comparing the two series, very small changes are observed. The central tendency of the data is preserved. A reduction in the extreme values and less variability is seen

after imputation. Although a shift in the values of skewness and kurtosis is seen, the distribution remains moderately left skewed and flat. The results from both the original and imputed time series indicate that neither of them follows a normal distribution according to the Anderson-Darling test. Fig. 4 shows the line plots of the time series, highlighting the outliers and imputed points, and also provides the LOESS trend plots.




Fig. 4. Annual CO2-E-LUC per capita time series plots with outliers and imputed points. LOESS trend of the original and imputed time series.

3.2 Non-linearity and Stationarity

3.2.1 Before First Differencing

Based on the results obtained from the BDS test (Tables 4-5) applied to the imputed time series of CO₂-E-LUC per capita, the standard normal statistics values are high and the *p*-values are less than 0.001. This occurs for all values of epsilon and embedding dimensions considered. Therefore, the initial hypothesis of the test is rejected. This fact indicates the presence of non-linearity or dependence in the time series data.

Table 4. BDS results for embedding dimension = 2, before and after first differencing.

First Differencing	Epsilon	Standard Normal	<i>p</i> -value
Before	0.5622	47.256	<0.001
	1.1244	44.122	<0.001
	1.6866	39.828	<0.001
	2.2489	34.212	<0.001
After	0.0932	8.0449	<0.001
	0.1864	6.269	<0.001
	0.2795	5.9787	<0.001
	0.3727	4.9951	<0.001

Table 5. BDS results for embedding dimension = 3, before and after first differencing.

First Differencing	Epsilon	Standard Normal (2)	Standard Normal (3)	<i>p</i> -value (2)	<i>p</i> -value (3)
Before	0.5622	46.125	72.716	<0.001	<0.001
	1.1244	42.704	51.523	<0.001	<0.001
	1.6866	38.775	42.586	<0.001	<0.001
	2.2489	33.572	34.694	<0.001	<0.001
After	0.0932	8.0288	10.3137	<0.001	<0.001
	0.1864	6.2542	7.5684	<0.001	<0.001
	0.2795	5.9498	6.9286	<0.001	<0.001
	0.3727	4.9703	5.6501	<0.001	<0.001

As can be seen in Table 6, non-linearity is confirmed also by the Teräsvirta Neural Network test, where the *p*-value is 4.961e-06 (<0.05). To further continue testing stationarity, we use tests that can handle non-linearity.

Table 6. Teräsvirta Neural Network test result, before and after first differencing.

First Differencing	Statistic		
	X-squared	Df	<i>p</i> -value
Before	24.428	2	4.961e-06
After	12.538	2	0.001894

The results of the KPSS test are displayed in Table 7. Across both short (4 lags) and long (13 lags) configurations for the KPSS test, the test statistics exceed the critical values at 5%, for stationarity around the mean (“mu”) and trend (“tau”). For the “mu” test, the short lag test statistic is 2.25, and the long lag test statistic is 0.8653. In the “tau” test, the short lag test statistic is 0.4681, and the long lag test statistic is 0.1873. This means that the null hypothesis of the test is rejected, and the imputed time series of CO₂-E-LUC per capita is neither level nor trend-stationary.

Table 8 provides the results of Zivot-Andrews’s test applied for varying lag values, from 1 to 40. Before first differencing, the Zivot-Andrew’s test provides the following results. In the “intercept” model, the test statistic -4.4048 is greater than all the critical values considered. This suggests that the series is non-stationary, even considering potential structural breaks. A break is identified at observation 141, and the optimal lag based on AIC is 18, indicating that the model explains a significant amount of autocorrelation in the data.

In the “trend” model, the test statistic -4.6469 is greater than the critical value at 1% but less than the critical value at 5% and 10%. According to this, there is weak evidence that the series remains non-stationary to a trend, and the potential structural break is detected at observation 126. The optimal lag is again 18, signifying similar autocorrelation patterns as in the “intercept” model.

In the “both” (intercept and trend) model, the test statistic of -6.0411 is less than all critical values (10%, 5%, and 1%). This suggests that the series is stationary when accounting for structural breaks in intercept and the trend. The structural break is identified at observation 90, and the optimal lag remains at 18, emphasizing once more long-term dependencies.

The data in Table 9 provides the results of the Breitung test. In both the models considered, we see that the value of the test statistic before differencing is greater than the critical values at all significance levels. This means that the null hypothesis of the test cannot be rejected.

In summary, after applying the three tests to check the stationarity of the imputed time series for CO2-E-LUC per capita, we come to the same conclusion: the series is non-stationary.

Table 7. KPSS test results on the imputed time series before and after first differencing.

Test Type	Lag	First Differencing	Test Statistic	Critical Values			
				(10%)	(5%)	(2.5%)	(1%)
KPSS (mu)	Short (4)	Before	2.25	0.347	0.463	0.574	0.739
		After	0.2407				
	Long (13)	Before	0.8653				
		After	0.1635				
KPSS (tau)	Short (4)	Before	0.4681	0.119	0.146	0.176	0.216
		After	0.0792				
	Long (13)	Before	0.1873				
		After	0.0556				

3.2.2 After First Differencing

To make the series stationary, we take the first differences. Then, we re-check if the series is non-linear. It turns out from the BDS test (Table 4 and Table 5) that the p-values remain less than 0.001, thus rejecting the null hypothesis of the test. It is noted that the values of standard normal statistics have decreased significantly compared to those before taking

the first differences. However, based on this test, the series remains non-linear.

The same phenomenon is observed in the Teräsvirta Neural Network test’s output (Table 6). Although the value of the test statistic has decreased, the p-value (0.001894) provides statistical evidence at the 0.05 significant level that the time series exhibits non-linearity.

Table 8. Zivot-Andrews Unit root test results before and after first differencing, with optimal lag based on AIC

Model Type	First Differencing	Test Statistic	Critical Values			Potential Break Point	Optimal Lag (AIC)
			10%	5%	1%		
Intercept	Before	-4.4048	-4.58	-4.80	-5.34	141	18
	After	-10.4096				108	1
Trend	Before	-4.6469	-4.11	-4.42	-4.93	126	18
	After	-9.214				130	1
Both	Before	-6.0411	-4.82	-5.08	-5.57	90	18
	After	-10.842				108	1

The first differencing has some effect on reducing non-linearity, but it is insufficient to fully eliminate it. Regarding the stationarity of the differenced time series, we continue testing with the same group of tests as before the first differencing due to the same conditions related to non-linearity.

Based on the results of the tests, we see that the first differencing has shifted the non-stationary series into a stationary one. The

KPSS test statistics (see Table 7) for both short and long lags are found to be less than the critical values. For the “mu” test, the short lag test statistic dropped to 0.2407, and the long lag test statistic fell to 0.1635, both less than the critical thresholds, indicating stationary around the mean. For the “tau” test, the short lag test statistic is 0.0792, and the long lag test statistic is 0.0556, providing evidence for trend stationarity after differencing.

Table 9. Breitung test results on the imputed time series before and after first differencing.

Test Type	Model	Lag	Test Statistic	Critical Values		
				(10%)	(5%)	(1%)
P-test	constant	1	Before: 29.112	4.33	3.17	1.91
			After: 0.4701			
P-test	trend	1	Before: 36.8543	6.86	5.66	4.05
			After: 1.4403			

After first differencing, the Zivot-Andrews’s test results (Table 8), present statistical evidence that the series becomes stationary across all model types. As a result, the null hypothesis of a unit root is rejected. In the “intercept” model, the test statistic value (-10.4096) is less than all critical values. The structural break shifts to observation 108, and the optimal lag based on AIC reduces to 1. In the “trend” model, the test statistic is -9.214, again less than all critical values, indicating that the series is now trend-stationary. The structural break occurs at observation 130, and the optimal lag is reduced to 1. In the “both” (intercept and trend) model, the test statistic of -10.842, less than all the critical values considered, confirms stationarity in both the intercept and trend after differencing. The structural break remains at 108, and the optimal lag is 1.

We see that after the first differencing, the autocorrelation is largely resolved, reducing the optimal lag to 1 in all three models, based on the Zivot-Andrews’s test results. Structural breaks still exist but occur at different points. While differencing eliminates non-stationarity, it does not entirely remove the impact of key structural changes in the series.

Results of the Breitung test after first differencing (Table 9) prove that the test statistic’s value becomes less than the critical values, in both models. In this case, the time series is stationary around a constant mean and also in a deterministic trend.

Taking first differences has successfully removed the unit root from the time series, making it suitable for further time series analysis or modeling. Our results related to non-linearity and stationarity of CO2 emissions data are consistent with other studies [41-43].

4. Conclusion

The statistical analysis of Albania's annual CO2 emissions from land-use change per capita was the main focus of this research. The Isolation Forest technique was employed to identify the outliers in the time series data and the LSTM model for imputation. The exploratory data analysis preserved the basic characteristics of the initial data distribution after imputation.

The BDS test and the Teräsvirta Neural Network test confirm the presence of non-linearity in the time series data, both before and after applying the first differences. The findings from the KPSS, Zivot-Andrews Unit Root, and Breitung tests indicate that the original time series was non-stationary, but first differencing transformed it into a stationary series, although non-linearity persisted.

The significance of this research lies in its detailed investigation of CO2-E-LUC per capita time series data in Albania, focusing on assessing the presence of non-linearity and stationarity. Furthermore, our study sheds light on the dynamic behavior of CO2 emissions from land-use changes. This is crucial for policymakers and researchers aiming to develop accurate predictive models and implement effective GHG emission reduction strategies. The study's limitations are rooted in the non-primary nature of the data we analyze, potentially affecting our conclusions. Future research on this dataset will employ advanced modeling techniques that account for non-linearity and stationarity, aiming for improved accuracy and reliability.

References

[1] Kamberi M, Islami B, Diku A, Profka D, Selfo G. National Greenhouse Gas Inventory Report under the Albania's First

- Biennial Update Report to UNFCCC. Ministry of Tourism and Environment. Tirana (AL): Ministry of Tourism and Environment; 2021. Available from: <https://www.undp.org/albania/publications/albanias-national-greenhouse-gas-inventory-report>
- [2] Raihan A, Begum RA, Nizam M, Said M, Pereira JJ. Dynamic impacts of energy use, agricultural land expansion, and deforestation on CO₂ emissions in Malaysia. *Environ Ecol Stat.* 2022;29:477-507.
- [3] IPCC. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Pachauri PK, Meyer LA, editors. Geneva: IPCC; 2014.
- [4] Friedlingstein P, Jones MW, O'Sullivan M, Andrew RM, Hauck J, Peters GP, et al. Global Carbon Budget 2019. *Earth Syst Sci Data.* 2019;11(4):1783-834. Available from: <https://essd.copernicus.org/articles-11/1783/2019/>
- [5] Jansen LJ, Carrai G, Petri M. Land-Use Change at Cadastral Parcel Level in Albania. In: Koomen E, Stillwell J, Bakema A, Scholten HJ, editors. *Modelling Land-Use Change*. Dordrecht: Springer; 2007. p. 25-44.
- [6] Qin Z, Zhu Y, Canadell JG, Chen M, Li T, Mishra U, Yuan W. Global spatially explicit carbon emissions from land-use change over the past six decades (1961–2020). *One Earth.* 2024;7:835-47.
- [7] Knez S, Štrbac S, Podbregar I. Climate change in the Western Balkans and EU Green Deal: status, mitigation and challenges. *Energy Sustain Soc.* 2022;12(1).
- [8] Baris Tuzemen O, Lyhagen J. Revisiting the role of climate change on crop production: evidence from Mediterranean countries. *Environ Dev Sustain.* 2024.
- [9] Pongratz J, Schwingshackl C, Bultan S, Obermeier W, Havermann F, Guo S. Land use effects on climate: current state, recent progress, and emerging topics. *Curr Clim Change Rep.* 2021;7:99-120.
- [10] Azam M, Adeleye BN. Impact of carbon emissions sources on life expectancy in Asia and the Pacific region. *Nat Resour Forum.* 2023;48(1):35-57.
- [11] Raihan A. The interrelationship amid carbon emissions, tourism, economy, and energy use in Brazil. *Carbon Res.* 2024;3(11).
- [12] Mitić P, Fedajev A, Radulescu M, Rehman A. The relationship between CO₂ emissions, economic growth, available energy, and employment in SEE countries. *Environ Sci Pollut Res.* 2022;30:16140-55.
- [13] Nicholls ZR, Gieseke R, Lewis J, Nauels A, Meinshausen M. Implications of nonlinearities between cumulative CO₂ emissions and CO₂-induced warming for assessing the remaining carbon budget. *Environ Res Lett.* 2020;15:074017.
- [14] European Commission. COMMISSION STAFF WORKING DOCUMENT: Albania 2023 Report. SWD(2023) 690 final, European Commission Council of the European Union; Brussels; 2023 [cited 2024]. Available from: <https://data-consilium.europa.eu/doc/document/ST-15098-2023-INIT/en/pdf>
- [15] Ritchie H, Roser M. CO₂ and greenhouse gas emissions. 2023 [cited 2024 Jun 07]. Available from: <https://github.com/owid-co2-data>
- [16] Ritchie H. CO₂ emissions dataset: our sources and methods. *Our World in Data.* 2022 [cited 2024 Jun 11]. Available from: <https://ourworldindata.org/co2-dataset-sources>
- [17] Brock WA, Dechert WD, Scheinkman JA, LeBaron B. A test for independence based on the correlation dimension. *Econom Rev.* 1996;15(3):197-235.

- [18] Cromwell JB, Labys WC, Terraza M. Testing for Linear or Nonlinear Dependence. In: *Univariate Tests for Time Series Models*. Thousand Oaks (CA): SAGE Publications, Inc.; 1994. p. 37-49.
- [19] Trapletti A, Hornik K, LeBaron B. *tseries*: Time Series Analysis and Computational Finance. R package version 0.10-56, CRAN (Comprehensive R Archive Network). 2024. Available from: <https://cran.r-project.org/web/packages/tseries/index.html>
- [20] Teräsvirta T. Specification, estimation, and evaluation of smooth transition autoregressive models. *J Am Stat Assoc*. 1994;89(425):208-18.
- [21] Teräsvirta T, Lin CF, Granger CW. Power of the Neural Network Linearity Test. *J Time Ser Anal*. 1993;14(2):209-20.
- [22] Di Narzo AF, Aznarte JL, Matthieu S. *tsDyn*: Nonlinear Time Series Models with Regime Switching. R package version 11.0.4.1, CRAN (Comprehensive R Archive Network). 2024. Available from: <https://cran.r-project.org/web/packages/tsDyn/index.html>
- [23] Kwiatkowski D, Phillips PC, Schmidt P, Shin Y. Testing the null hypothesis of stationarity against the alternative of a unit root: How sure are we that economic time series have a unit root? *J Econom*. 1992;54(1-3):159-78.
- [24] Pfaff B, Zivot E, Stigler M. *urca*: Unit Root and Cointegration Tests for Time Series Data. R package version 1.3-4, CRAN (Comprehensive R Archive Network). 2024. Available from: <https://cran.r-project.org/web/packages/urca/index.html>
- [25] Zivot E, Andrews DW. Further evidence on the Great Crash, the oil-price shock, and the unit-root hypothesis. *J Bus Econ Stat*. 1992;10(3):251-70.
- [26] Elliott G, Rothenberg T, Stock J. Efficient tests for an autoregressive unit root. *Econometrica*. 1996;64(4):813-36.
- [27] MacKinnon JG. Critical values for cointegration tests. In: Engle RF, Granger CW, editors. *Long-Run Economic Relationships: Readings in Cointegration*. Oxford (UK): Oxford University Press; 1991. p. 267-76.
- [28] Akaike H. A new look at the statistical model identification. *IEEE Trans Autom Control*. 1974;19(6):716-23.
- [29] Cortes D. *Isolation-Based Outlier Detection*. R package Version 0.6.1-1, CRAN (Comprehensive R Archive Network). 2024. Available from: <https://github.com/david-cortes/isotree>
- [30] Liu FT, Ting KM, Zhou ZH. Isolation forest. In: *Proc. 8th IEEE Int. Conf. Data Mining (ICDM)*; Pisa, Italy; 2008. p. 413–22.
- [31] Hochreiter S, Schmidhuber J. Long short-term memory. *Neural Comput*. 1997;9(8):1735-80.
- [32] Hyndman RJ, Koehler AB. Another look at measures of forecast accuracy. *Int J Forecast*. 2006;22(4):679-88.
- [33] Allaire J, Chollet F. *keras*: R Interface to Keras. R package version 2.15.0, CRAN. 2024. Available from: <https://github.com/rstudio/keras/tree/r2>
- [34] Allaire J, Tang Y. *tensorflow*: R Interface to TensorFlow. R package version 2.16.0. 2024. Available from: <https://github.com/rstudio/tensorflow>
- [35] Hamner B, Frasco M. *Evaluation Metrics for Machine Learning*. R package version 0.1.4, CRAN. 2018. Available from: <https://github.com/mfrasco/Metrics>
- [36] Hyndman R, Athanasopoulos G, Bergmeir C, Caceres G, Chhay L, Kuroptev K, et al. *Forecasting Functions for Time Series and Linear Models*. R package version 8.23.0, CRAN. 2024. Available from: <https://github.com/robjhyndman/forecast>

- [37] Anderson TW, Darling DA. Asymptotic theory of certain "goodness of fit" criteria based on stochastic processes. *Ann Math Stat.* 1952;23(2):193-212.
- [38] R Core Team. *loess: Local Polynomial Regression Fitting*. 2024. Available from: <https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/loess>
- [39] Cleveland WS, Grosse E, Shyu WM. Local regression models. In: Chambers JM, Hastie TJ, editors. *Statistical Models in S*. Pacific Grove (CA): Wadsworth & Brooks/Cole; n.d. Chapter 8.
- [40] R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna (AT): R Foundation for Statistical Computing; 2023. Available from: <https://www.R-project.org/>
- [41] Solarin SA, Gil-Alana LA, Lafuente C. Persistence in carbon footprint emissions: an overview of 92 countries. *Carbon Manag.* 2019;10(4):405-15.
- [42] Sohail A, Du J, Abbasi BN, Ahmed Z. The nonlinearity and nonlinear convergence of CO2 emissions: Evidence from top highest emitting countries. *Environ Sci Pollut Res.* 2022;29:59466-82.
- [43] Churchill SA, Inekwe J, Ivanovski K, Smyth R. Stationarity properties of per capita CO2 emissions in the OECD in the very long-run: A replication and extension analysis. *Energy Econ.* 2020;90:104868.